

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS

S0

elev8™ film coated tablets
Sceletium tortuosum (L.) N.E. Br. (Kanna).
Sugar free.

COMPLEMENTARY MEDICINE: OTHER HERBAL MEDICINES

This unregistered medicine has not been evaluated by the SAHPRA for its quality, safety or intended use.

Read all of this leaflet carefully because it contains important information for you.

elev8™ is available without a doctor's prescription, for you to maintain your health. Nevertheless, you still need to use elev8™ carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your healthcare provider or pharmacist if you need more information or advice.

What is in this leaflet

1. What elev8™ is and what it is used for.
2. What you need to know before you take elev8™.
3. How to take elev8™.
4. Possible side effects.
5. How to store elev8™.
6. Contents of the pack and other information.

1. What elev8™ is and what it is used for

elev8™ assists in coping with stress, irritability, emotional strain, mood fluctuations and might support cognitive function in adults.

2. What you need to know before you take elev8™

Do not take elev8™:

- if you are hypersensitive (allergic) to *Sceletium tortuosum* (L.) N.E. Br. or any of the other ingredients of elev8™ (see **What elev8™ contains**);
- if you are taking psychiatric medicine, including medicine to stabilise your moods, antidepressants or anti-anxiety medicine.

Warnings and precautions

None known.

Other medicines and elev8™

Always tell your healthcare provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

elev8™ with food, drink and alcohol

See section 3 (**How to take elev8™**).

Pregnancy, breastfeeding and fertility

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other healthcare provider for advice before taking this medicine. Safety and efficacy of elev8™ during pregnancy and breastfeeding have not been established. Do not use this medicine if you are pregnant or breastfeeding your baby.

Driving and using machines

elev8™ might make you feel drowsy. Do not drive a vehicle, operate machinery, or do anything else that requires your attention until you know how elev8™ affects you.

3. How to take elev8™

Do not share your medicines with any other person.

Always take elev8™ exactly as described in this leaflet, or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The usual dose is:

Adults and children from 12 years of age: 1 – 2 tablets in the morning with a glass of water.

If you have the impression that the effect of elev8™ is too strong or too weak, tell your doctor or pharmacist.

Do not exceed the recommended dosage.

elev8™ can be taken with or without food.

elev8™ should not be taken with alcohol.

If you take more elev8™ than you should

In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison centre.

Take this leaflet and the rest of the remaining elev8™ with you so the doctor will know what you have taken.

If you forget to take elev8™

- Take the missed dose as soon as you remember.
- Do not take a double dose to make up for the forgotten individual doses.

4. Possible side effects

elev8™ can have side effects.

Not all side effects reported for elev8™ are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking elev8™, please consult your healthcare provider for advice.

If any of the following happens, stop taking elev8™ and tell your doctor immediately or go to the casualty department at your nearest hospital:

- swelling of your hands, feet, ankles, face, lips, mouth or throat, which may cause difficulty in swallowing or breathing;
- rash or itching;
- fainting.

These are all very serious side effects. If you have them, you may have had a serious allergic reaction to elev8™. You may need urgent medical attention or hospitalisation.

Tell your doctor if you notice any of the following:

Side effects occurring less frequently:

- loss of appetite;
- headaches;
- lack of energy or feeling tired;
- stimulation;
- sedation;
- mild diarrhoea (runny, loose stool), mild stomach cramps.

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor or pharmacist. You can also report side effects to SAHPRA via the “6.04 Adverse Drug Reaction Reporting Form”, found online **under SAHPRA’s publications**:

<https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of elev8™.

5. How to store elev8™

- Store all medicines out of reach of children.
- Store at or below 25 °C.
- Store the blister strips in the carton.
- Protect from light and moisture.

- Do not store in the bathroom.
- Do not use after the expiry date printed on the carton or blister strip.
- Return all unused medicine to your pharmacist.
- Do not dispose of unused medicine in drains and sewerage systems (e.g. toilets).

6. Contents of the pack and other information

What elev8™ contains

Each film coated tablet contains 25 mg Zembrin®, equivalent to:

Sceletium tortuosum (L.) N.E. Br. (Kanna) 50 mg
[above ground herb, providing 100 µg mesembrine-type alkaloids]

The other ingredients are calcium phosphate dibasic dihydrate, coating [containing castor oil (E1503), polyvinyl alcohol-polyethylene glycol-*graft*-copolymer, Quinoline Yellow aluminium lake (E104), shellac (E904), talcum, titanium dioxide (E172)], croscarmellose sodium (E468), magnesium stearate (E572), microcrystalline cellulose (E460 (i)), polyvinyl pyrrolidone (E1201), vinylpyrrolidone-vinyl acetate copolymer.

What elev8™ looks like and contents of the pack

Yellow, round, convex, unscored film coated tablet.

PVDC/aluminium blister strip containing 10 tablets, packed in boxes of 10, 30, 60 and 90.

Holder of Certificate of Registration

Brunel Laboratoria (Pty) Ltd

1 Van Tonder Street

Sunderland Ridge

Centurion

0157

info@brunel.co.za

This leaflet was last revised in

Will be allocated by SAHPRA upon registration.

Registration number

Will be allocated by SAHPRA upon registration.

Access to the corresponding Professional Information

The Professional Information for elev8™ is available on <https://www.brunel.co.za/elev8/> or can be requested from the contact details above.

PASIËNTINLIGTINGSBLAD

SKEDULERINGSSTATUS

S0

elev8™ filmbedekte tablette
Sceletium tortuosum (L.) N.E. Br. (Kanna).
Suikervry.

KOMPLEMENTÊRE MEDISYNE: ANDER KRUIEMEDISYNE

Hierdie ongeregistreerde medisyne is nie deur die SAHPRA geëvalueer vir sy gehalte, veiligheid of beoogde gebruik nie.

Lees hierdie hele inligtingsblad noukeurig deur want dit bevat inligting wat belangrik is vir jou.

elev8™ is beskikbaar sonder 'n dokter se voorskrif, vir jou om jou gesondheid te onderhou. Nietemin, moet jy elev8™ versigtig gebruik om die beste resultate daaruit te kry.

- Hou hierdie inligtingsblad. Dit mag nodig wees dat jy dit weer lees.
- Vra gerus jou gesondheidsorgverskaffer of apteker indien jy verdere inligting of advies nodig het.

Wat is in hierdie inligtingsblad

1. Wat elev8™ is, en waarvoor dit gebruik word.
2. Wat jy moet weet voordat jy elev8™ neem.
3. Hoe om elev8™ te neem.
4. Moontlike nuwe-effekte.
5. Hoe om elev8™ te bêre.
6. Inhoud van die verpakking en ander inligting.

1. Wat elev8™ is, en waarvoor dit gebruik word

elev8™ help om stres, prikkelbaarheid, emosionele spanning en gemoedskommeling te hanteer en mag kognitiewe funksie in volwassenes ondersteun.

2. Wat jy moet weet voordat jy elev8™ neem

Moet nie elev8™ neem:

- indien jy hipersensitief (allergies) is vir *Sceletium tortuosum* (L.) N.E. Br. of vir enige van die ander bestanddele van elev8™ nie (sien **Wat elev8™ bevat**);
- indien jy psigiatryse medisyne neem, insluitende medisyne om jou gemoedstoestand te stabiliseer, antidepressante of medisyne vir angstigheid.

Waarskuwings en voorsorgmaatreëls

Geen bekend nie.

Ander medisyne en elev8™

Sê altyd vir jou gesondheidsorgverskaffer indien jy enige ander medisyne neem. (Dit sluit alle komplementêre of tradisionele medisyne in.)

elev8™ met eet-, drinkgoed en alkohol

Sien afdeling 3 (**Hoe om elev8™ te neem**).

Swangerskap, borsvoeding en vrugbaarheid

Indien jy swanger is of jou baba borsvoed, dink dat jy swanger mag wees of beplan om swanger te raak, raadpleeg asseblief jou dokter, apteker of ander gesondheidsorgverskaffer voordat jy die medisyne neem. Die veiligheid en effektiwiteit van elev8™ tydens swangerskap en borsvoeding is nie vasgestel nie. Moet nie hierdie medisyne gebruik as jy swanger is of jou baba borsvoed nie.

Bestuur en die gebruik van masjiene

elev8™ mag jou lomerig laat voel. Moet nie 'n voertuig bestuur, masjinerie hanteer, of enigiets anders doen wat jou aandag vereis totdat jy weet hoe elev8™ jou affekteer nie.

3. Hoe om elev8™ te neem

Moet nie jou medisyne met enige ander persoon deel nie.

Neem elev8™ altyd presies soos aangedui in hierdie inligtingsblad, of soos jou dokter of apteker vir jou verduidelik het. Vra gerus jou dokter of apteker indien jy onseker is.

Die gewone dosis is:

Volwassenes and kinders vanaf 12-jarige ouderdom: 1 – 2 tablette soggens, met 'n glas water.

Indien jy die indruk kry dat die uitwerking van elev8™ te sterk of te swak is, bespreek dit met jou dokter of apteker.

Moet nie die aanbevole dosis oorskry nie.

elev8™ kan met of sonder kos geneem word.

elev8™ moet nie saam met alkohol geneem word nie.

As jy meer elev8™ neem as wat jy moet

In die geval van oordosering, raadpleeg jou dokter of apteker. Indien nie een van hulle beskikbaar is nie, kontak die naaste hospitaal of gifsentrum.

Neem hierdie inligtingsblad en die oorblywende elev8™ saam met jou sodat die dokter kan sien wat jy geneem het.

As jy vergeet om elev8™ te neem

- Neem die dosis wat jy oorgeslaan het sodra jy onthou.
- Moet nie 'n dubbele dosis neem om vir die vergete individuele dosis op te maak nie.

4. Moontlike nuwe-effekte

elev8™ kan nuwe-effekte hê.

Nie alle nuwe-effekte wat vir elev8™ gerapporteer is, word in hierdie inligtingsblad ingesluit nie. Indien jou algemene gesondheid verswak of indien jy enige ongunstige effekte ondervind terwyl jy elev8™ neem, raadpleeg asseblief jou gesondheidsorgverskaffer.

Indien enige van die volgende gebeur, hou op om elev8™ te neem en sê dadelik vir jou dokter, of gaan na die ongevalle-afdeling van jou naaste hospitaal:

- swelling van jou hande, voete, enkels, gesig, lippe, mond of keel, wat probleme kan veroorsaak met sluk en asemhaling;
- uitslag of gejeuk;
- floute.

Hierdie is baie ernstige nuwe-effekte. Indien jy enige daarvan ondervind, mag jy 'n ernstige allergiese reaksie op elev8™ gehad het. Jy mag dringend mediese bystand of hospitalisasie benodig.

Sê vir jou dokter indien jy enige van die volgende opmerk:

Nuwe-effekte wat minder gereeld voorkom:

- aptytverlies;
- hoofpyn;
- gebrek aan energie of gevoel van moegheid;
- stimulasie;
- sedasie;
- geringe diaree (lopende, los stoelgang), ligte maagkrampe.

Indien jy enige nuwe-effekte opmerk wat nie in hierdie inligtingsblad genoem word nie, sê asseblief vir jou dokter of apteker daarvan.

Aanmelding van nuwe-effekte

Indien jy nuwe-effekte ondervind, bespreek dit met jou dokter of apteker. Jy kan ook nuwe-effekte aan SAHPRA rapporteer via die “**6.04 Adverse Drug Reaction Reporting Form**”, wat aanlyn by SAHPRA se publikasies beskikbaar is: <http://www.sahpra.org.za/Publications/Index/8>. Deur nuwe-effekte te rapporteer kan jy help om meer inligting rakende die veiligheid van elev8™ te verskaf.

5. Hoe om elev8™ te bêre

- Bêre alle medisyne buite die bereik van kinders.
- Bêre teen of onder 25 °C.
- Stoor die stulpstrokies in die karton.
- Beskerm teen direkte sonlig en vog.

- Moet nie in die badkamer stoor nie.
- Moet nie gebruik word na die vervaldatum wat op die karton of stulpstrokie gedruk is nie.
- Neem alle ongebruikte medisyne terug na jou apteker.
- Moet nie ongebruikte medisyne in dreine en rioolstelsels (bv. toilette) weggooi nie.

6. Inhoud van die verpakking en ander inligting

Wat elev8™ bevat

Elke filmbedekte tablet bevat 25 mg Zembrin®, gelykstaande aan:

Sceletium tortuosum (L.) N.E. Br. (Kanna) 50 mg
[bogronde dele van kruie, lewer 100 µg mesembrien-tipe alkalioiede]

Die ander bestanddele is kalsiumfosfaat dibasiese dihidraat, filmlagie [bevattende kasterolie (E1503), polivinielalkohol-poliëtileenglikol-gehegte-kopolimeer, Quinoline Yellow aluminiumlak (E104), skilferlak (E904), talkum, titaandioksied, (E172)], natriumkroskarmellose (E468), magnesiumstearaat (E572), mikrokristallyne sellulose (E460 (i)), polivinielpirrolidoon (E1201), vinielpirrolidoon-vinielasetaat-kopolimeer.

Hoe elev8™ lyk en die inhoud van die verpakking

Geel, ronde, konvekse, filmbedekte tablet sonder 'n middellyn.

PVDC/aluminium stulpstrook wat 10 tablette bevat, verpak in 'n buitenste karton met 10, 30, 60 of 90 tablette.

Houer van die Registrasiesertifikaat

Brunel Laboratoria (Edms) Bpk
Van Tonderstraat 1
Sunderland Ridge
Centurion
0157
info@brunel.co.za

Hierdie inligtingsblad is hersien in

Sal met registrasie deur SAHPRA aangedui word.

Registrasienommer

Sal met registrasie deur SAHPRA toegeken word.

Toegang tot die ooreenstemmende Professionele Inligtingsblad

Die Professionele Inligtingsblad vir elev8™ is beskikbaar by <https://www.brunel.co.za/elev8/> of dit kan aangevra word deur die kontakbesonderhede hier bo te gebruik.